

DRUG FACTS

Active ingredient(s)		Purpose
Use(s)		
<p>Warnings</p> <p>Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:</p> <ul style="list-style-type: none">• hives• facial swelling• asthma (wheezing)• shock• skin reddening• rash• blisters <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you:</p> <ul style="list-style-type: none">• are age 60 or older• have had stomach ulcers or bleeding problems• take a blood thinning (anticoagulant) or steroid drug• take other drugs containing NSAID (aspirin, ibuprofen, naproxen, or others)• have 3 or more alcoholic drinks every day while using this product• take more or for a longer time than directed <p>Do not use</p> <ul style="list-style-type: none">• if you have ever had an allergic reaction to any other pain reliever/fever reducer• right before or after heart surgery <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none">• problems or serious side effects from taking pain relievers or fever reducers• stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain• ulcers• bleeding problems• high blood pressure• heart or kidney disease• taken a diuretic• reached age 60 or older <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none">• taking aspirin for heart attack or stroke because ibuprofen may decrease this benefit of aspirin• taking any other drug containing an NSAID (prescription or nonprescription)• taking a blood thinning (anticoagulant) or steroid drug• under a doctor's care for any serious condition• taking any other drug <p>When using this product</p> <ul style="list-style-type: none">• take with food or milk if stomach upset occurs• long term continuous use ay increase the risk of heart attack or stroke <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none">• you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.• pain gets worse or lasts more than 10 days• fever gets worse or lasts more than 3 days• stomach pain or upset gets worse or lasts• redness or swelling is present in the painful area• any new symptoms appear <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.</p>		
Do not use		
Ask a doctor before use if you have		
Ask a doctor or pharmacist before use if you are		
When using this product		
Stop use and ask a doctor if		
If pregnant or breast-feeding		
Keep out of reach of children		
Directions		
<p>Other information</p> <p>*Compared to the active ingredient of Advil®</p> <p>Drug Facts</p> <ul style="list-style-type: none">• temporarily relieves minor aches and pains due to:• headache• muscular aches• minor pain or arthritis• toothache• backache• the common cold• menstrual cramps• temporarily reduces fever• do not take more than directed• the smallest effective dose should be used• do not take longer than 10 days, unless directed by a doctor (see Warnings)		
Adults and children 12 years and older		<ul style="list-style-type: none">• take 1 tablet every 4 to 6 hours while symptoms persist• if pain or fever does not respond to 1 tablet, 2 tablets may be used• do not exceed 6 tablets in a 24-hour period

Children under 12 years

• ask a doctor

Distributed by:
Amneal Pharmaceuticals
Glasgow, KY 42141

*Amneal Pharmaceuticals is not affiliated with the owner of the trademark Advil®
Rev. 10-2008

Inactive ingredients